

DAFTAR PUSTAKA

- [1] S. A. Malik, O. P. Singh, A. Nurifhan, and M. B. Malarvili, “Portable Respiratory CO₂ Monitoring Device for Early Screening of Asthma,” no. December, 2016, doi: 10.15224/978-1-63248-113-9-61.
- [2] M. Farooqui *et al.*, “A Non-invasive device and automated monitoring system using peak flow meter for asthma patients,” *ICCAIS 2020 - 3rd Int. Conf. Comput. Appl. Inf. Secur.*, 2020, doi: 10.1109/ICCAIS48893.2020.9096881.
- [3] T. T. Kean and M. B. Malarvili, “Analysis of capnography for asthmatic patient,” *ICSIPIA09 - 2009 IEEE Int. Conf. Signal Image Process. Appl. Conf. Proc.*, pp. 464–467, 2009, doi: 10.1109/ICSIPIA.2009.5478699.
- [4] C. S. Hong, “Development of electronic respiratory for detecting asthma,” no. June, p. 19, 2017.
- [5] fran hegarty Kabita shakya, catherine deegan, “DETERMINATION FREQUENCY ETCO2.pdf,” vol. 4, no. 2, p. 56, 2003, doi: 10.21427/D73J0N.
- [6] AAST, “Technical Guideline End-Tidal CO₂,” vol. 2, no. 7. American Association of Sleep Technologists, pp. 1–19, 2018.
- [7] Resironics Inc. and R. Inc., “Capnography Reference Handbook.” pp. 1–36, 2004, [Online]. Available: http://oem.resironics.com/wp/CapnographyReferenceHandbook_OEM1220A.pdf.
- [8] M. Khezri, “Capnography,” no. July, 2015, doi:

10.13140/RG.2.1.4701.5523.

- [9] C. V. Egleston, H. Ben Aslam, and M. A. Lambert, “Capnography for monitoring non-intubated spontaneously breathing patients in an emergency room setting,” *Emerg. Med. J.*, vol. 14, no. 4, pp. 222–224, 1997, doi: 10.1136/emj.14.4.222.
- [10] L. Safitrin, “Rancang Bangun Sistem Pendekripsi kadar CO₂ Hasil Ekspirasi,” 2010.
- [11] H. Aminiahidashti, S. Shafiee, A. Z. Kiasari, and M. Sazgar, “Applications of End-Tidal Carbon Dioxide (ETCO 2) Monitoring in Emergency Department; a Narrative Review,” vol. 6, no. 1, pp. 1–6, 2018.
- [12] F. J. Cereceda-sánchez, “Capnography as a tool to detect metabolic changes in patients cared for in the emergency setting,” 2017, doi: 10.1590/1518-8345.1756.2885.
- [13] J. A.-H. Ahmad O. Muhtaseb, Firas M, “Design and implementation of portable diabetic ketoacidosis detector,” *Foreign Aff.*, pp. 1–76, 2015, doi: 10.1017/CBO9781107415324.004.
- [14] mb malarvif Singh, Om Prakash, “Review of Infrared Carbon-Dioxide Sensors and Capnogram Features for Developing Asthma-Monitoring Device,” vol. 12, pp. 1–6, 2018, doi: 10.7860/JCDR/2018/35870.12099.
- [15] T. Jo *et al.*, “Usefulness of measurement of end-tidal CO₂ using a portable capnometer in patients with chronic respiratory failure receiving long-term oxygen therapy,” *Intern. Med.*, vol. 59, no. 14, pp. 1711–1720, 2020, doi: 10.2169/internalmedicine.3320-19.

- [16] B. You, R. Peslin, C. Duvivier, V. D. Vu, and J. P. Grilliat, “Expiratory capnography in asthma: Evaluation of various shape indices,” *Eur. Respir. J.*, vol. 7, no. 2, pp. 318–323, 1994, doi: 10.1183/09031936.94.07020318.
- [17] D. Santoso and F. D. Setiaji, “Design and Implementation of Capnograph for Laparoscopic Surgery,” *Int. J. Inf. Electron. Eng.*, vol. 3, no. 5, pp. 523–528, 2013, doi: 10.7763/ijiee.2013.v3.370.
- [18] S. J. Capistrano, “(12) Patent Application Publication (10) Pub . No .: US 2015 / 0238722 A1,” 2015.
- [19] K. L. Chan, M. T. V. Chan, and T. Gin, “Mainstream vs. sidestream capnometry for prediction of arterial carbon dioxide tension during supine craniotomy,” *Anaesthesia*, vol. 58, no. 2, pp. 149–155, 2003, doi: 10.1046/j.1365-2044.2003.03035.x.
- [20] M. W. Tjokorda Gde Agung Senopathi, “The End Tidal CO₂ correlation with decreased cardiac output measured by ultrasonic cardiac output monitor in intubated ICU patients,” vol. 6, p. 5, 2017.
- [21] J. G. Goepp, “EFFORT-INDEPENDENT, PORTABLE, USER-OPERATED CAPNOGRAPH DEVICES AND RELATED METHODS,” US2009/0118632A1, 2009.
- [22] M. G. Srabanti, “Design of NDIR based CO₂ sensor to detect human respiratory CO₂,” *Sustain.*, vol. 11, no. 1, pp. 1–14, 2019.
- [23] C. Pantazopoulos, T. Xanthos, I. Pantazopoulos, A. Papalois, E. Kouskouni, and N. Iacovidou, “A Review of Carbon Dioxide Monitoring During Adult

Cardiopulmonary Resuscitation,” *A Rev. Carbon Dioxide Monit. Dur. Adult Cardiopulm. Resusc.*, vol. 24, no. 11, pp. 1053–1061, 2015, doi: 10.1016/j.hlc.2015.05.013.

- [24] M. Kazemi and M. Malarvili, “Analysis of Capnogram Using Linear Predictive Coding (LPC) to Differentiate Asthmatic Conditions,” *J. Tissue Sci. Eng.*, vol. 02, no. 05, pp. 2–5, 2012, doi: 10.4172/2157-7552.1000111.
- [25] C. C. Rasera and P. M. Gewehr, “Association between Capnogram and Respiratory Flow Rate Waveforms during Invasive Mechanical Ventilation,” *Assoc. between capnogram Respir. flow rate waveforms Dur. invasive Mech. Vent.*, vol. 3, no. 2, pp. 80–84, 2013, doi: 10.7763/ijbbb.2013.v3.169.
- [26] O. P. Singh and M. B. Malarvili, “Review of infrared carbon-dioxide sensors and capnogram features for developing asthma-monitoring device,” *J. Clin. Diagnostic Res.*, vol. 12, no. 10, pp. OE01–OE06, 2018, doi: 10.7860/JCDR/2018/35870.12099.