

DAFTAR PUSTAKA

- [1] I. A. E. Widiastuti, R. Cholidah, G. W. Buanayuda, and I. B. Alit, “Deteksi Dini Faktor Risiko Penyakit Kardiovaskuler pada Pegawai Rektorat Universitas Mataram,” *J. Pengabdi. Magister Pendidik. IPA*, vol. 4, no. 1, 2021, doi: 10.29303/jpmipi.v4i1.604.
- [2] J. Misbach, “Stroke di Indonesia : a fisrt large prospective hospital based study of acute stroke in 28 hospitals in Indonesia,” *J. Clin. Neurosci.*, vol. 8, no. 3, pp. 245–249, 2009.
- [3] O. S. Anita Shinta Kusuma, “Penerapan Prosedur Latihan Range of Motion (Rom) Pasif Sedini Mungkin Pada Pasien Stroke Non Hemoragik (Snh),” *Syntax Lit.*, vol. 5, no. 10, pp. 1015–1021, 2020, [Online]. Available: <https://www.jurnal.syntaxliterate.co.id/index.php/syntax-literate/article/view/1706/1614>.
- [4] A. Hamzah and Sugiyanto, “Strengthening of Health Locus of Control could Increase the Independence of Post Stroke Patients in Implementing the Daily Activities at Home,” *J.*

Nurs. Care, vol. 03, no. 02, 2014, doi: 10.4172/2167-1168.1000152.

- [5] C. M. Rismawanti, W. Natalya, and I. Isytiaroh, “Penerapan Latihan Range Of Motion (ROM) aktif pada pasien pasca stroke di Desa Sampangan Pekalongan,” *Pros. Semin. Nas. Kesehat.*, vol. 1, pp. 1350–1354, 2021, doi: 10.48144/prosiding.v1i.834.
- [6] I. Jahra Alhamid, N. Norma, and O. Lopulalan, “ANALISIS FAKTOR RISIKO TERHADAP KEJADIAN STROKE,” *Nurs. Arts*, vol. 12, no. 2, pp. 100–109, Dec. 2018, doi: 10.36741/jna.v12i2.82.
- [7] P. Lindsay, K. L. Furie, S. M. Davis, G. A. Donnan, and B. Norrving, “World stroke organization global stroke services guidelines and action plan,” *Int. J. Stroke*, vol. 9, no. A100, pp. 4–13, 2014, doi: 10.1111/ijs.12371.
- [8] K. L. Lanctôt *et al.*, “Canadian Stroke Best Practice Recommendations: Mood, Cognition and Fatigue following Stroke, 6th edition update 2019,” *Int. J. Stroke*, vol. 15, no. 6, pp. 668–688, 2020, doi: 10.1177/1747493019847334.

- [9] D. Rofifah, “FAKTOR RESIKO TERHADAP KEJADIAN STROKE Di RSUD Dr. KOESNADI BONDOWOSO,” *Pap. Knowl. . Towar. a Media Hist. Doc.*, pp. 12–26, 2020.
- [10] M. Hughes and G. Y. H. Lip, “Stroke and thromboembolism in atrial fibrillation: A systematic review of stroke risk factors, risk stratification schema and cost effectiveness data,” *Thromb. Haemost.*, vol. 99, no. 2, pp. 295–304, 2008, doi: 10.1160/TH07-08-0508.
- [11] B. Stegmayr and K. Asplund, “Diabetes as a risk factor for stroke. A population perspective,” *Diabetologia*, vol. 38, no. 9, pp. 1061–1068, 1995, doi: 10.1007/s001250050392.
- [12] A. K. Boehme, C. Esenwa, and M. S. V. Elkind, “Stroke Risk Factors, Genetics, and Prevention,” *Circ. Res.*, vol. 120, no. 3, pp. 472–495, 2017, doi: 10.1161/CIRCRESAHA.116.308398.
- [13] H. P. Kuntono, “Penatalaksanaan Fisioterapi Pada Kondisi Osteoarthritis,” *Temu Ilmiyah IFI, Kediri*, 2005.

- [14] X. D. Xu, Y. Jin, Y. Zhao, W. W. Wu, and X. D. Wang, “Orthotic effect of functional electrical stimulation on the improvement of walking in stroke patients with foot drop: A systematic review,” *Chinese J. Evidence-Based Med.*, vol. 13, no. 6, pp. 735–740, 2013, doi: 10.7507/1672-2531.20130130.
- [15] R. Setiawan, S. Sabar, and M. Madi, “Design of An Active Power Ankle-Foot Orthosis with Myoelectric Control for Drop-Foot Rehabilitation,” *Int. J. Electr. Energy Power Syst. Eng.*, vol. 4, no. 2, pp. 139–144, 2021, doi: 10.31258/ijeepe.4.2.139-144.
- [16] K. Dunning, M. W. O’Dell, P. Kluding, and K. McBride, “Peroneal Stimulation for Foot Drop after Stroke,” *Am. J. Phys. Med. Rehabil.*, vol. 94, no. 8, pp. 649–664, 2015, doi: 10.1097/PHM.0000000000000308.
- [17] A. E. K. Sari, J. Purwono, and Ludiana, “Penerapan Range of Motion (Rom) Pasif Untuk Meningkatkan Kekuatan Otot Pasien Pasca Stroke Di Wilayah Kerja Puskesmas Purwosari Kec .

Metro Utara Application of the Passive Range of Motion (Rom) To Improve Muscular Strength of Patients After Stroke in,” *J. Cendekia Muda*, vol. 1, no. 2, pp. 209–215, 2021.

- [18] V. T. Huttami and A. C. Hidajah, “the Utilization of Golden Period of Ischemic Stroke in Patients in Productive Ages,” *Indones. J. Public Heal.*, vol. 15, no. 3, p. 258, 2020, doi: 10.20473/ijph.v15i3.2020.258-265.
- [19] R. Ambarika and N. A. Anggraini, “Golden Period in Terms of Knowledge Family of Early Detection of Stroke for Stroke Patients in Installation of Emergency,” *Malaysian J. Med. Res.*, vol. 06, no. 03, pp. 30–34, 2022, doi: 10.31674/mjmr.2022.v6i03.004.
- [20] C. V. Oramas, D. D. Keluarga, and C. V. Oramas, “RANCANG BANGUN SISTEM PEREKAMAN SINYAL EMG UNTUK MONITORING PERKEMBANGAN PASIEN PASCA STROKE,” no. Bagian I, p. 2016, 2016.
- [21] A. G. Bharatkumar, K. E. Daigle, M. G. Pandy, Q. Cai, and J. K. Aggarwal, “Lower limb kinematics

- of human walking with the medial axis transformation,” *Motion Non-Rigid Articul. Objects Work. Proc.*, pp. 70–76, 1994, doi: 10.1109/mnrao.1994.346252.
- [22] K. Szymańska, M. Domżalski, and A. Stolarczyk, “Differences in male and female lower limb joint kinematics during walking,” *Rehabil. Med.*, vol. 23, no. 3, pp. 39–46, 2019, doi: 10.5604/01.3001.0013.3767.
- [23] N. R. Agusrianto, “Penerapan Latihan Range of Motion (Rom) Pasif terhadap Peningkatan Kekuatan Otot Ekstremitas pada Pasien dengan Kasus Stroke,” *J. Ilm. Kesehat.*, vol. 2, no. 2, pp. 61–66, 2020.
- [24] B. D. M. Chaparro-Rico, D. Cafolla, P. Tortola, and G. Galardi, “Assessing stiffness, joint torque and ROM for paretic and non-paretic lower limbs during the subacute phase of stroke using lokomat tools,” *Appl. Sci.*, vol. 10, no. 18, 2020, doi: 10.3390/APP10186168.
- [25] A. Laudanski, B. Brouwer, and Q. Li, “Measurement of lower limb joint kinematics using inertial sensors during stair ascent and

descent in healthy older adults and stroke survivors,” *J. Healthc. Eng.*, vol. 4, no. 4, pp. 555–576, 2013, doi: 10.1260/2040-2295.4.4.555.

- [26] J. L. Stephenson, S. J. De Serres, and A. Lamontagne, “The effect of arm movements on the lower limb during gait after a stroke,” *Gait Posture*, vol. 31, no. 1, pp. 109–115, 2010, doi: 10.1016/j.gaitpost.2009.09.008.
- [27] F. Y. Liang, F. Gao, J. Cao, S. W. Law, and W. H. Liao, “Gait Synergy Analysis and Modeling on Amputees and Stroke Patients for Lower Limb Assistive Devices,” *Sensors*, vol. 22, no. 13, 2022, doi: 10.3390/s22134814.
- [28] N. F. SARI, “PENATALAKSANAAN FISIOTERAPI PADA KASUS POST DISLOKASI ELBOW DEXTRA DI RSUD PANEMBAHAN SENOPATI BANTUL,” pp. 1–13, 2012.
- [29] G. Vagenas and B. Hoshizaki, “Functional Asymmetries and Lateral Dominance in the Lower Limbs of Distance Runners,” *Int. J. Sport Biomech.*, vol. 7, no. 4, pp. 311–329, 2016, doi:

- 10.1123/ijsb.7.4.311.
- [30] M. Zuk and M. Trzeciak, “Anatomical protocol for gait analysis: Joint kinematics measurement and its repeatability,” *J. Theor. Appl. Mech.*, vol. 55, no. 1, pp. 369–376, 2017, doi: 10.15632/jtam-pl.55.1.369.
 - [31] R. J. Butler, S. Marchesi, T. Royer, and I. S. Davis, “The Effect of a Subject-Specific Amount of Lateral Wedge on Knee,” *J. Orthop. Res. Sept.*, vol. 25, no. June, pp. 1121–1127, 2007, doi: 10.1002/jor.
 - [32] D. Yuliansyah, “Deteksi Kelelahan Otot Menggunakan Sinyal Emg Dan Detektor Gaya Pada Gerak Dasar Ekstensi Dan Fleksi Knee-Joint Untuk Evaluasi Penggunaan Functional Electrical Stimulation Pada Sistem Rehabilitasi Lower Limb,” *Sepuluh Nop. Inst. Technol.*, p. 171, 2017.
 - [33] K. Takeda, G. Tanino, and H. Miyasaka, “Review of devices used in neuromuscular electrical stimulation for stroke rehabilitation,” *Med. Devices Evid. Res.*, vol. 10, pp. 207–213, 2017, doi: 10.2147/MDER.S123464.
 - [34] S. K. Sabut, C. Sikdar, R. Kumar, and M.

Mahadevappa, “Functional electrical stimulation of dorsiflexor muscle: Effects on dorsiflexor strength, plantarflexor spasticity, and motor recovery in stroke patients,” *NeuroRehabilitation*, vol. 29, no. 4, pp. 393–400, 2011, doi: 10.3233/NRE-2011-0717.

- [35] T. Yan, C. W. Y. Hui-Chan, and L. S. W. Li, “Functional electrical stimulation improves motor recovery of the lower extremity and walking ability of subjects with first acute stroke: A randomized placebo-controlled trial,” *Stroke*, vol. 36, no. 1, pp. 80–85, 2005, doi: 10.1161/01.STR.0000149623.24906.63.
- [36] C. T. Freeman, A. M. Hughes, J. H. Burridge, P. H. Chappell, P. L. Lewin, and E. Rogers, “A model of the upper extremity using FES for stroke rehabilitation,” *J. Biomech. Eng.*, vol. 131, no. 3, 2009, doi: 10.1115/1.3005332.
- [37] P. M. Kluding *et al.*, “Foot drop stimulation versus ankle foot orthosis after stroke: 30-week outcomes,” *Stroke*, vol. 44, no. 6, pp. 1660–1669, 2013, doi: 10.1161/STROKEAHA.111.000334.

- [38] J. T. Mesin, F. T. Industri, and U. I. Indonesia, “PERANCANGAN SISTEM KENDALI ALAT BANTU Pernyataan keaslian,” 2021.
- [39] D. Shi, W. Zhang, W. Zhang, and X. Ding, “A Review on Lower Limb Rehabilitation Exoskeleton Robots,” *Chinese J. Mech. Eng.*, 2019, doi: 10.1186/s10033-019-0389-8.
- [40] M. M, “Kontraksi Otot Skelet,” *J. MensSana*, vol. 2, no. 2, p. 69, 2017, doi: 10.24036/jm.v2i2.25.
- [41] ADELIA RACHMAWATI, “PENGEMBANGAN STANDAR OPERASIONAL PROSEDUR (SOP) TEKNIK DISTRAKSI DENGAN TERAPI MUROTTAL TERHADAP PENURUNAN INTENSITAS NYERI PADA PASIEN FRAKTUR POST ORIF,” *Pap. Knowl. . Towar. a Media Hist. Doc.*, vol. 3, no. April, pp. 49–58, 2015.
- [42] ANNISA’USH SHOLIHAH, “PENGARUH MOBILISASI DINI TERHADAP PENINGKATAN KEKUATAN OTOT PADA PASIEN PASCA STROKE ISKEMIK DI RSUD DR. HARJONO PONOROGO,” no. December, 2017.

- [43] 1689–1699.
<https://doi.org/10.1017/CBO9781107415324.004>
Francisco, A. R. L. (2013). Journal of Chemical Information and Modeling, 53(9), “Anatomi dan Fisiologi Otot Rangka,” *J. Chem. Inf. Model.*, vol. 53, no. 9, pp. 1689–1699, 2013.
- [44] W. Kamelia1, “PENGARUH LATIHAN ROM (RANGE OF MOTION) TERHADAP KEKUATAN OTOT PADA PASIEN POST STROKE The Effect Of Rom (Range Of Motion) Exercise On Muscle Strength In Post Stroke Patients yang bersifat fungsional , sedangkan pada pasien pasca stroke akan mengalami,” vol. 2, no. 2, pp. 109–114, 2021.
- [45] K. I. Nur Rahayu, “Pengaruh pemberian latihan,” *J. Keperawatan PENGARUH PEMBERIAN Latih. RANGE MOTION TERHADAP Kemamp. Mot. PADA PASIEN POST STROKE DI RSUD GAMBIRAN*, vol. 6, no. 2010, pp. 102–107, 2015, [Online]. Available: <https://media.neliti.com/media/publications/138683-ID-pengaruh-pemberian-latihan-range-of->

moti.pdf.

- [46] U. Sarah, S. Bambang, and K. Bm Wara, “Pengaruh Latihan Range of Motion (Rom) Terhadap Fleksibilitas Sendi Lutut Pada Lansia Di Panti Wreda Wening Wardoyo Ungaran,” *Pengaruh Latih. ROM Media Ners*, vol. 1, no. 2, p. 49, 2007.
- [47] R. Setiawan, H. H. Triharminto, and M. Fahrurrozi, “Gesture Control Menggunakan IMU MPU 6050 Metode Kalman Filter Sebagai Kendali Quadcopter,” *Pros. Semin. Nas. Sains Teknol. dan Inov. Indones.*, vol. 3, no. November, pp. 411–422, 2021, doi: 10.54706/senastindo.v3.2021.133.
- [48] Mpu6050, *DataSheet MPU6050*. .
- [49] M. S. Rosyidi, “Rancang Bangun Alat Pembersih Dan Penyortir Ukuran Telur Asin Berbasis Arduino Mega 2560,” *J. Skripsi*, pp. 1–17, 2015.
- [50] M. Suari, “Pemanfaatan Arduino nano dalam Perancangan Media Pembelajaran Fisika,” *Nat. Sci. J.*, vol. 3, no. 1, pp. 474–480, 2017, [Online]. Available: www.ecadio.com.
- [51] A. Rahmadiansyah, E. Orlanda, M. Wijaya, H. W. Nugroho, and R. Firmansyah, “Perancangan

Sistem Telemetri Untuk Mengukur Intensitas Cahaya Berbasis Sensor Light Dependent Resistor Dan Arduino Uno,” *JEEE-U (Journal Electr. Electron. Eng.*, vol. 1, no. 1, pp. 15–21, 2017, doi: 10.21070/jeee-u.v1i1.760.

